Supporting Information

Efficient Bioconjugation of 5-Fluoro-5-Deoxy-Ribose (FDR) to RGD Peptides for Positron Emission Tomography (PET) Imaging of $\alpha_{\nu}\beta_{3}$ Integrin Receptor

Sergio Dall'Angelo,^a Qingzhi Zhang,^b Ian Fleming,^c Monica Piras,^a Lutz F. Schweiger,^a David O' Hagan,^{b,*} Matteo Zanda^{a,*}

¹John Mallard Scottish PET Centre, School of Medical Sciences, University of Aberdeen, Foresterhill, Aberdeen, AB25 2ZD.

²EastChem School of Chemistry and biological Sciences research Complex, University of St Andrews, North Haugh, St Andrews, KY16 9ST, UK.

³Aberdeen Biomedical Imaging Centre, Lilian Sutton Building, Foresterhill, Aberdeen AB25 2ZD, UK




Figure S1 ¹H NMR of (cRGDfC)-Ahm in D₂O with solvent suppression





Figure S2 2D TOCSY spectrum of (cRGDfC)-Ahm in D_2O












Figure S5 ¹H NMR of (cRGDfC)-Ahm-FDR in D₂O with solvent suppression






















Figure S12 ^{19}F NMR of c(RGDfK)-Aoa-FDR in D_2O

