Macrocycle synthesis by trimerization of boronic acids around a hexaol template, and recognition of polyols by resulting macrocyclic oligoboronic acids

Dennis Stoltenberg and Ulrich Lüning

Figure 1	¹³ C-NMR of triester 13a
Figure 2	¹³ C-NMR of triester 13b
Figure 3	¹³ C-NMR of triester 13c
Figure 4	¹³ C-NMR of triester 13d
Figure 5	¹³ C-NMR of triester 14a
Figure 6	¹³ C-NMR of triester 14b
Figure 7	¹³ C-NMR of triester 16
Figure 8	¹³ C-NMR of triester 17

Figure 1. ¹³C-NMR of triester 13a




Figure 2. ¹³C-NMR of triester 13b




Figure 3. ¹³C-NMR of triester 13c




Figure 4. ¹³C-NMR of triester 13d




Figure 5. ¹³C-NMR of triester 14a

 $X = (CH_2)_n CH = CH(CH_2)_n$

14a: *n* = 3




Figure 6. ¹³C-NMR of triester 14b

$$X = (CH_2)_n CH = CH(CH_2)_n$$

14b: *n* = 4




Figure 7. ¹³C-NMR of triester 16




Figure 8. ¹³C-NMR of triester 17

