

Progress towards water-soluble triazole-based selective MMP-2 inhibitors

*Benjamin Fabre,^a Kamila Filipiak,^{a,b} Jose María Zapico,^a Natalia Díaz,^c Rodrigo J. Carbajo,^d Anne K. Schott,^d M. Paz Martínez-Alcázar,^a Dimas Suárez,^c Antonio Pineda-Lucena,^d Ana Ramos,^{*a} Beatriz de Pascual-Teresa^{*a}*

^aDepartamento de Química, Facultad de Farmacia, Universidad CEU San Pablo,
Urbanización Montepríncipe, 28668 Madrid, Spain.

^bDepartment of Molecular Biology, Faculty of Mathematics and Natural Sciences, The
John Paul II Catholic University of Lublin, 20-718, Lublin, Poland

^cDepartamento de Química Física y Analítica, Universidad de Oviedo, C/ Julián
Clavería, 8. 33006 Oviedo, Asturias, Spain

^dStructural Biology Laboratory, Medicinal Chemistry Department, Centro de
Investigación Príncipe Felipe, C/ Eduardo Primo Yúfera 3, 46012 Valencia, Spain

SUPPORTING INFORMATION

Table of Content

Time evolution of the MMP-2 backbone atoms RMSD of the MD trajectory.....	S2
¹H and ¹³C NMR spectra of 1e-i and 13a-j.....	S2

Time evolution of the MMP-2 backbone atoms RMSD of the MD trajectory










Figure S1. Time evolution of the Root-Mean-Square Deviation (RMSD, in Å) for the protein backbone atoms (except the *N*-term coil) of the MD trajectory.



^1H and ^{13}C NMR spectra of 1e-i and 13a-j



































13f



13f

