Supplementary Information for:

Synthesis of saxitoxin derivatives bearing guanidine and urea groups at C13 and evaluation of their inhibitory activity on voltage-gated sodium channel

Takafumi Akimoto,<sup>a</sup> Asako Masuda,<sup>a</sup> Mari Yotsu-Yamashita,<sup>b</sup> Takatsugu Hirokawa<sup>c</sup> and Kazuo Nagasawa<sup>\*a</sup>

[a] Department of Biotechnology and Life Science,

Faculty of Technology, Tokyo University of Agriculture and Technology

2-24-16 Nakamachi, Koganei, Tokyo, 184-8588, Japan

Fax: (+81) 42-388-7295

E-mail: knaga@cc.tuat.ac.jp

\_\_\_\_\_

[b] Graduate of Agriculture Science, Tohoku University

1-1 Tsutsumidori-Amamiyamachi, Aoda-ku, Sendai, Miyagi, 981-8555, Japan

[c] Computational Biology Research Center National Institute of Advanced Industrial Science and Technology

2-4-7, Aomi, Koto-ku, Tokyo, 135-0064, Japan.

## Table of Contents:

1. <sup>1</sup> H NMR and <sup>13</sup> C NMR spectra	
---	--

1. <sup>1</sup>H NMR and <sup>13</sup>C NMR spectra







Electronic Supplementary Material (ESI) for Organic & Biomolecular Chemistry This journal is The Royal Society of Chemistry 2013

















Electronic Supplementary Material (ESI) for Organic & Biomolecular Chemistry This journal is The Royal Society of Chemistry 2013

















