

Supporting Information

Synthesis and anionophoric activities of dimeric polyamine-sterol conjugates: the impact of rigid vs flexible linkers

Yong-Ming Lu,^a Li-Qun Deng,^a Xi Huang,^a Jin-Xiang Chen,^a Bo Wang,^b Zhong-Zhen Zhou,^a Guan-Song Hu^a and Wen-Hua Chen

*. ^a

^a *School of Pharmaceutical Sciences, Southern Medical University, Guangzhou 510515, P. R. China*

^b *School of Chemistry and Chemical Engineering, Sun Yat-Sen University, Guangzhou 510275, P. R. China*

Figures S1~6. ¹H NMR spectra of compounds **1-4** and boc-protected **2**; and ¹³C NMR spectrum of compound **2**.




Figure S1. ¹H NMR spectrum (400 MHz, CD₃OD) of compound **3**.




Figure S2. ¹H NMR spectrum (400 MHz, CD₃OD) of compound 4.




Figure S3. ¹H NMR spectrum (400 MHz, CD₃OD) of compound **1**.




Figure S4. ^1H NMR spectrum (400 MHz, CD_3OD) of boc-protected **2**.





Figure S5. ¹H NMR spectrum (400 MHz, CD₃OD) of compound **2**.



Figures S6. ^{13}C NMR spectrum (100 MHz, CD_3OD) of compounds **2**.