

Supporting Information

for

**Electrochemical behaviour of new dimeric esters and amides
derived from caffeic acid in dimethylsulfoxide**

Analilia Sánchez, Omar Martínez-Mora, Evelin Martínez-Benavidez,
Javier Hernández, Zaira Domínguez* and Magali Salas-Reyes*

Unidad de Servicios de Apoyo en Resolución Analítica, Universidad Veracruzana,
Apdo. Postal 575, Xalapa, Ver., México

* Corresponding authors.

E-mail addresses: magasalas@uv.mx (Magali Salas-Reyes), zdominguez@uv.mx (Zaira
Domínguez)

Table of contents

1. Cyclic voltammetry of compound 6 at different scan rates.....	S1
2. Cyclic voltammetry of compound 8 at different scan rates.....	S2
3. ¹³ C NMR spectrum of compound 6	S3
4. ¹³ C NMR spectrum of compound 7	S4
5. ¹³ C NMR spectrum of compound 8	S5
6. ¹³ C NMR spectrum of compound 9	S6




Figure S1. Cyclic voltammetry of compound **6** at different scan rates (from 0.025 to 0.5 V s⁻¹). Insert: dependence of the anodic peak current I on the square root of the potential scan rate (v).




Figure S2. Cyclic voltammetry of compound **8** at different scan rates (from 0.05 to 0.8 V s⁻¹). Insert: dependence of the cathodic peak current **IIIa** and **IIIb** on the square root of the potential scan rate (v).




Figure S3. ¹³C NMR spectrum of compound **6** (100.5 MHz; d₆-DMSO).




Figure S4. ^{13}C NMR spectrum of compound 7 (100.5 MHz; d_6 -DMSO).




Figure S5. ¹³C NMR spectrum of compound **8** (75.5 MHz; CD₃OD).




Figure S6. ¹³C NMR spectrum of compound 9 (75.5 MHz; CD₃OD).