

Electronic Supplementary Information

3-NO₂-5,10,15-triarylcorrolato-Cu as a versatile platform for synthesis of novel 3-functionalized corrole derivatives

Manuela Stefanelli,^a Marco Mancini,^a Michele Raggio,^a Sara Nardis,^{*a} Frank R. Fronczek,^b Gregory McCandless,^{b,c} Kevin M. Smith,^b and Roberto Paolesse^a

nardis@scienze.uniroma2.it;

^a *Dipartimento di Scienze e Tecnologie Chimiche, Università di Roma Tor Vergata, via della Ricerca Scientifica 1, 00133 Rome, Italy*

^b *Department of Chemistry, Louisiana State University, Baton Rouge, LA 70803, USA*

^c *Department of Chemistry, The University of Texas at Dallas, Richardson, TX 75080, USA*

Copies of ¹H NMR and IR spectra




Figure S1: ^1H NMR spectrum of 3-NHC(O)CH₃-TtBuCorrCu(2)




Figure S2: ^1H NMR spectrum of 3-[NHC(O)(CH₂)₄CH₃]-TtBuCorrCu(3)




Figure S3: ^1H NMR spectrum of 3-NHC(O)CH₂Cl-TtBuCorrCu(4)




Figure S4: ^1H NMR spectrum of 3-NHC(O)CH₂N₃-TtBuCorrCu(5)




Figure S5: IR spectrum of 3-NHC(O)CH₂N₃-TtBuCorrCu(5)




Figure S6: ¹H NMR spectrum of 3-NHC(O)CH₂(4-tolyl-triazole)TtBuCorrCu(6)




Figure S6: ^1H NMR spectrum of 3-NHC(O)H-TtBuCorrCu(7)