Supporting information for

Selective cleavage of the Cα-Cβ linkage in lignin model compounds via Baeyer-Villiger oxidation

Nikhil D. Patil,^a Soledad G. Yao,^a Mark S. Meier,^{*,a} Justin K. Mobley^{a,b} and Mark Crocker,^{a,b}

^aDepartment of Chemistry, University of Kentucky, Lexington, KY 40506, USA ^bCenter for Applied Energy Research, Lexington, KY 40511, USA

NMR Spectra

1.	Compound 2	page S2
2.	Compound 3	page S4
3.	Compound 4	page S5
4.	Compound 5	page S5
5.	Compound 6	page S6
6.	Compound 8	page S7
7.	Compound 15	page S8
8.	Compound 18	page S9
9.	Compound 19	page S10
10.	Compound 26	page S11
11.	Compound 27	page S12
12.	Compound 28	page S13
13.	Compound 29	page S14
14.	Compound 30	page S15
15.	Compound 31	page S16
16.	Compound 33	page S17
17.	Compound 34	page S18
18.	Compound 35	page S19
19.	Compound 36	page S20
20.	Compound 38	page S21



NMR spectra were measured at 400 MHz (¹H) and 100 MHz (¹³C) in CDCl₃ solution. ¹H NMR and ¹³C NMR spectra of 2



















S7



S8









S11













¹H NMR of compound **31**















