Electronic Supplementary Material (ESI) for Organic & Biomolecular Chemistry. This journal is © The Royal Society of Chemistry 2014

SUPPORTING INFORMATION OF COMPOUNDS (5a-aa)

Synthesis and biological evaluation of imidazo[1,5-a]pyridine-benzimidazole hybrids as inhibitors of both tubulin polymerization and PI3K/Akt pathway

Ahmed Kamal, a, b* A.V. Subba Rao, V. Lakshma Nayak, N.V. Subba Reddy, a

Konderu Swapna, ^b G. Ramakrishna ^a and Mallika Alvala ^b

^aMedicinal Chemistry and Pharmacology, CSIR-Indian Institute of Chemical Technology, Hyderabad, 500 007, India

^bDepartment of Medicinal Chemistry, National Institute of Pharmaceutical Education and Research (NIPER), Hyderabad, 5000 037, India

Experimental

Chemistry


Materials and Methods.

¹H NMR spectra were recorded on Avance 300, Inova 400, Avance 500, and Bruker 600 MHz spectrometers using tetramethylsilane (TMS) as the internal standard. Chemical shifts are reported in parts per million (ppm) downfield from tetramethylsilane. Spin multiplicities are described as s (singlet), brs (broad singlet), d (doublet), dd (double doublet), t (triplet), q (quartet), and or m (multiplet). Coupling constants are reported in Hertz (Hz). Melting points were determined in a capillary tube using an Electrothermal apparatus (Model IA9200) and are uncorrected. The IR spectra were recorded by employing a Nicolet FTIR model MX-1 spectrophotometer. Analytical thin layer chromatography (TLC) was performed on MERCK precoated silica gel 60- F254 (0.5 mm) glass plates. Visualization of the spots on TLC plates was achieved either by exposure to iodine vapour or UV light or by dipping the plates into β-naphthol or to ethanolic anisaldehyde or to ethanolic ninhydrin solution and heating the plates to 120 °C.


Column chromatography was performed using silica gel 60-120 or 100-200 mesh. Moisture sensitive reactions were carried out using standard syringe septum techniques and under inert atmosphere of nitrogen. All solvents and reagents were purified by standard techniques. *O*-Phenylenediamines (11a-i) and substituted benzoyl chlorides (7a-c) are commercially available and used as received. All evaporation of solvents was carried out under reduced pressure on rotary evaporator below 45 °C. The names of all the compounds given in the experimental section were taken from Chem Ultra, Version 11.0.

General procedure for synthesis of compounds (5a-aa):


To a solution appropriate aldehyde (10a-c) in ethanol (10 mL), appropriate *o*-phenylenediamines (11a-c) (1 mmol) and sodium metabisulpite (2 mmol) dissolved in H₂O (5 mL) were added sequentially and then reaction mixture was refluxed for 4 h. After completion of reaction, solvent was removed by vacuo to give crude product which was extracted with ethyl acetate. The combined organic extracts were dried over anhydrous Na₂SO₄ and concentrated to give crude products which were further purified by column chromatography by using ethylacetate and hexane as solvent system to afford pure compounds (5a-aa).




¹H NMR SPECTRA OF COMPOUND 5a




¹³C NMR SPECTRA OF COMPOUND 5a




HIGH RESOLUTION MASS SPECTRA OF COMPOUND 5a




¹H NMR SPECTRA OF COMPOUND 5b




¹³C NMR SPECTRA OF COMPOUND 5b




HIGH RESOLUTION MASS SPECTRA OF COMPOUND 5b




¹H NMR SPECTRA OF COMPOUND 5C




¹³C NMR SPECTRA OF COMPOUND 5c




HIGH RESOLUTION MASS SPECTRA OF COMPOUND 5c




¹H NMR SPECTRA OF COMPOUND 5d




¹³C NMR SPECTRA OF COMPOUND 5d




HIGH RESOLUTION MASS SPECTRA OF COMPOUND 5d




¹H NMR SPECTRA OF COMPOUND 5d




¹³C NMR SPECTRA OF COMPOUND 5e




HIGH RESOLUTION MASS SPECTRA OF COMPOUND 5e




¹H NMR SPECTRA OF COMPOUND 5f




¹³C NMR SPECTRA OF COMPOUND 5f




HIGH RESOLUTION MASS SPECTRA OF COMPOUND 5f




¹H NMR SPECTRA OF COMPOUND 5g




HIGH RESOLUTION MASS SPECTRA OF COMPOUND 5g




¹H NMR SPECTRA OF COMPOUND 5h




¹³C NMR SPECTRA OF COMPOUND 5h




HIGH RESOLUTION MASS SPECTRA OF COMPOUND 5h




¹H NMR SPECTRA OF COMPOUND 5i




¹³C NMR SPECTRA OF COMPOUND 5i




HIGH RESOLUTION MASS SPECTRA OF COMPOUND 5i




 1 H NMR SPECTRA OF COMPOUND 5j




¹³C NMR SPECTRA OF COMPOUND 5j




HIGH RESOLUTION MASS SPECTRA OF COMPOUND 5j




¹H NMR SPECTRA OF COMPOUND 5k




¹³C NMR SPECTRA OF COMPOUND 5k




HIGH RESOLUTION MASS SPECTRA OF COMPOUND 5k




¹H NMR SPECTRA OF COMPOUND 51




¹³C NMR SPECTRA OF COMPOUND 51




HIGH RESOLUTION MASS SPECTRA OF COMPOUND 51




¹H NMR SPECTRA OF COMPOUND 5m




 $^{13}\mathrm{C}$ NMR SPECTRA OF COMPOUND 5m




HIGH RESOLUTION MASS SPECTRA OF COMPOUND 5m




¹H NMR SPECTRA OF COMPOUND 5n




¹³C NMR SPECTRA OF COMPOUND 5n




HIGH RESOLUTION MASS SPECTRA OF COMPOUND 5n




¹H NMR SPECTRA OF COMPOUND 50




¹³C NMR SPECTRA OF COMPOUND 50




HIGH RESOLUTION MASS SPECTRA OF COMPOUND 50




¹H NMR SPECTRA OF COMPOUND 5p




¹³C NMR SPECTRA OF COMPOUND 5p




HIGH RESOLUTION MASS SPECTRA OF COMPOUND 5p




¹H NMR SPECTRA OF COMPOUND 5q




¹³C NMR SPECTRA OF COMPOUND 5q




HIGH RESOLUTION MASS SPECTRA OF COMPOUND 5q




¹H NMR SPECTRA OF COMPOUND 5r




HIGH RESOLUTION MASS SPECTRA OF COMPOUND 5r




¹H NMR SPECTRA OF COMPOUND 5s




¹³C NMR SPECTRA OF COMPOUND 5s




HIGH RESOLUTION MASS SPECTRA OF COMPOUND 5s




¹H NMR SPECTRA OF COMPOUND 5t




¹³C NMR SPECTRA OF COMPOUND 5t




HIGH RESOLUTION MASS SPECTRA OF COMPOUND 5t




¹H NMR SPECTRA OF COMPOUND 5u




¹³C NMR SPECTRA OF COMPOUND 5u




HIGH RESOLUTION MASS SPECTRA OF COMPOUND 5u




¹H NMR SPECTRA OF COMPOUND 5v




¹³C NMR SPECTRA OF COMPOUND 5v




HIGH RESOLUTION MASS SPECTRA OF COMPOUND 5v




¹H NMR SPECTRA OF COMPOUND 5w




 $^{13}\mathrm{C}$ NMR SPECTRA OF COMPOUND 5w




HIGH RESOLUTION MASS SPECTRA OF COMPOUND 5w




¹H NMR SPECTRA OF COMPOUND 5x




¹³C NMR SPECTRA OF COMPOUND 5x




HIGH RESOLUTION MASS SPECTRA OF COMPOUND 5x




¹H NMR SPECTRA OF COMPOUND 5y




¹³C NMR SPECTRA OF COMPOUND 5y




HIGH RESOLUTION MASS SPECTRA OF COMPOUND 5y




¹H NMR SPECTRA OF COMPOUND 5z




HIGH RESOLUTION MASS SPECTRA OF COMPOUND 5z



¹H NMR SPECTRA OF COMPOUND 5aa



¹³C NMR SPECTRA OF COMPOUND 5aa



HIGH RESOLUTION MASS SPECTRA OF COMPOUND 5aa