

Photophysical and Laser Emission Studies of 8-Polyphenylene-Substituted BODIPY Dyes in Liquid Solution and in Solid Polymeric Matrices

M. Álvarez ^a, A. Costela ^{*}, and I. García-Moreno

Instituto de Química Física "Rocasolano", C.S.I.C., Serrano 119, 28006 Madrid, Spain.

F. Amat-Guerri and M. Liras¹

Instituto de Química Orgánica, C.S.I.C., Juan de la Cierva 3, 28006 Madrid, Spain.

R. Sastre

Instituto de Ciencia y Tecnología de Polímeros, C.S.I.C., Juan de la Cierva 3, 28006 Madrid, Spain.

F. López Arbeloa, J. Bañuelos Prieto, and I. López Arbeloa

Departamento de Química Física, UPV-EHU, Apartado 644, 48080 Bilbao, Spain


Supporting Information

Spectra and fluorescence decay curves of model (P2Ar1Ac and P3Ar1Ac) and monomer (P2Ar1MA and P3Ar1MA) dyes incorporated in PMMA and covalently linked to PMMA respectively, illustrating the absence of shift concerning the absorption band and the bathochromic shift of the emission band


^{*} Corresponding autor. Tel.: +34-91-561-9400; fax: + 34-91-564-2431
E-mail address: acostela@iqfr.csic.es (A. Costela).

^a Current address: Max Planck Institute for Polymer Research, Ackermannweg 10, D-55128 Mainz, Germany.

¹ Current address: Universidad Miguel Hernández, Dpto. de Ciencia y Tecnología de los Materiales, Avda. Ferrocarril, 03202 Elche, Alicante, Spain.



Absorption and emission (normalized to the fluorescence quantum yield) spectra (A) and fluorescence decay curves (B) of P2ArAc incorporated in PMMA (a) and P2ArMA covalently linked to PMMA (b). Dye concentration $0.8 \times 10^{-3} \text{ M}$ and sample thickness 0.5 mm.



Absorption and emission (normalized to the fluorescence quantum yield) spectra (A) and fluorescence decay curves (B) of P3ArAc incorporated in PMMA (a) and P3ArMA covalently linked to PMMA (b). Dye concentration $0.8 \times 10^{-3} \text{ M}$ and sample thickness 0.5 mm.