

A facile one-step synthesis to ionic liquids-based cross-linked polymeric nanoparticle and its
application for CO₂ fixation

Yubing Xiong,* Yujiao Wang, Hong Wang, Rongmin Wang

*Key Laboratory of Eco-Environment-Related Polymer Materials Ministry of Education, Key
Laboratory of Polymer Materials of Gansu Province, College of Chemistry and Chemical
Engineering, Northwest Normal University, Lanzhou 730070, China*

Corresponding author:

E-mail address: yubing_xiong@163.com (Y.B. Xiong)

Postal address: 967 East Anning Road, Northwest Normal University, Lanzhou 730070, China.

Tel.: +86-931-7970075; Fax: +86-931-7970358




Figure S1. ^1H NMR spectrum of 4-vinylbenzyl-tributylphosphorous chloride




Figure S2. ^{13}C NMR spectrum of 4-vinylbenzyl-tributylphosphorous chloride




Figure S3. Photographs of cross-linked polymer emulsions in different solvents. a) absolute methanol; b) methanol; c) absolute ethanol; d) ethanol; e) absolute isopropanol; f) isopropanol.




Figure S4. Photographs of cross-linked polymer emulsions with different molar ratio of EGMA to PIL




Figure S5 Photographs of cyclocaddition reaction mixture. (a) just after the reaction; (b) after toluene was added.




Figure S6




Figure S7




Figure S8




Figure S9




Figure S10




Figure S11 SEM and TEM images of used CLPN.