

Supporting Information

One-Step Preparation of Reduction-Responsive Poly(ethylene glycol)-Poly(amino acid)s Nanogels as Efficient Intracellular Drug Delivery Platform

Jianxun Ding,^{‡ac} Fenghua Shi,^{‡ab} Chunsheng Xiao,^{ac} Lin Lin,^a Li Chen,^{*b} Chaoliang He,^a Xiuli Zhuang,^{*a} Xuesi Chen^a

^aKey Laboratory of Polymer Ecomaterials, Changchun Institute of Applied Chemistry, Chinese Academy of Sciences, Changchun 130022, P. R. China.

^bDepartment of Chemistry, Northeast Normal University, Changchun 130022, P. R. China.

^cGraduate University of the Chinese Academy of Sciences, Beijing 100039, P. R. China.

[‡]These authors contributed equally to this paper.

Corresponding Author: Li Chen and Xiuli Zhuang

E-mail: chenl686@nenu.edu.cn and zhuangxl@ciac.jl.cn; Tel/Fax: +86-431-85262367.




Figure S1. ^1H NMR spectrum of mPEG-NH₂ (in CDCl₃).




Figure S2. Excitation spectra of pyrene in PBS of NGs at different concentration ($\lambda_{em} = 390 \text{ nm}$) (A), and the intensity ratio (I_{333}/I_{331}) as a function of concentration of various NGs (B).