

Electronic Supplementary Information

A New Route to Functional Polymers: Atom-Economical Synthesis of Poly(pyrazolynaphthalene)s by Rhodium-Catalyzed Oxidative Polycoupling of Phenylpyrazole and Internal Diynes

*Meng Gao,^a Jacky W. Y. Lam,^a Yajing Liu,^a Jie Li,^a and Ben Zhong Tang^{*abc}*

^a Department of Chemistry, Institute for Advanced Study, State Key Laboratory of Molecular Neuroscience, Institute of Molecular Functional Materials and Division of Biomedical Engineering, The Hong Kong University of Science & Technology (HKUST), Clear Water Bay, Kowloon, Hong Kong, China. E-mail: tangbenz@ust.hk

^b Guangdong Innovative Research Team, SCUT-HKUST Joint Research Laboratory, State Key Laboratory of Luminescent Materials and Devices, South China University of Technology (SCUT), Guangzhou, 510640, China

^c HKUST Shenzhen Research Institute, No. 9 Yuexing 1st RD, South Area, Hi-tech Park, Nanshan, Shenzhen, China 518057.

Table of Contents

Fig. S1 IR spectra of (A) **1**, (B) **2(4)** and (C) their polymer **P1/2(4)**.

Fig. S2 ¹H NMR spectra of CDCl₃ solutions of (A) **1**, (B) **2(6)**, (C) a mixture of **5–8** and (D) **P1/2(6)** (sample taken from Table 3, no. 2).

Fig. S3 ¹³C NMR spectra of CDCl₃ solutions of (A) **1**, (B) **2(6)**, (C) a mixture of **5–8** and (D) **P1/2(6)** (sample taken from Table 3, no. 2). The solvent peaks were marked with asterisks.

Fig. S4 ¹H NMR spectra of CDCl₃ solutions of (A) **1**, (B) **2(8)**, (C) a mixture of **5–8** and (D) **P1/2(8)** (sample taken from Table 3, no. 3).

Fig. S5 ¹³C NMR spectra of CDCl₃ solutions of (A) **1**, (B) **2(8)**, (C) a mixture of **5–8** and (D)

P1/2(8) (sample taken from Table 3, no. 3). The solvent peaks were marked with asterisks.

Fig. S6 ^1H NMR spectra of CDCl_3 solutions of (A) **1**, (B) **2(10)**, (C) a mixture of **5–8** and (D) P1/2(10) (sample taken from Table 3, no. 4).

Fig. S7 ^{13}C NMR spectra of CDCl_3 solutions of (A) **1**, (B) **2(10)**, (C) a mixture of **5–8** and (D) P1/2(10) (sample taken from Table 3, no. 4). The solvent peaks were marked with asterisks.

Fig. S8 ^1H NMR spectra of CDCl_3 solutions of (A) **1**, (B) **3** and (C) P1/3 (sample taken from Table 3, no. 5).

Fig. S9 ^{13}C NMR spectra of CDCl_3 solutions of (A) **1**, (B) **3** and (C) P1/3 (sample taken from Table 3, no. 5). The solvent peaks were marked with asterisks.

Fig. S10 Size distribution of nanoparticles of P1/3 suspended in 99% aqueous mixture. Concentration: 10 μM .

Fig. S11 Plot of quantum yield versus the solvent composition of THF/water mixture of P1/3.




Fig. S1 IR spectra of (A) **1**, (B) **2(4)** and (C) their polymer P1/2(4).




Fig. S2 ¹H NMR spectra of CDCl₃ solutions of (A) **1**, (B) **2(6)**, (C) a mixture of **5–8** and (D) **P1/2(6)** (sample taken from Table 3, no. 2).




Fig. S3 ¹³C NMR spectra of CDCl₃ solutions of (A) **1**, (B) **2(6)**, (C) a mixture of **5–8** and (D) **P1/2(6)** (sample taken from Table 3, no. 2). The solvent peaks were marked with asterisks.




Fig. S4 ¹H NMR spectra of CDCl₃ solutions of (A) **1**, (B) **2(8)**, (C) a mixture of **5–8** and (D) **P1/2(8)** (sample taken from Table 3, no. 4).




Fig. S5 ¹³C NMR spectra of CDCl₃ solutions of (A) **1**, (B) **2(8)**, (C) a mixture of **5–8** and (D) **P1/2(8)** (sample taken from Table 3, no. 3). The solvent peaks were marked with asterisks.




Fig. S6 ^1H NMR spectra of CDCl_3 solutions of (A) **1**, (B) **2(10)**, (C) a mixture of **5–8** and (D) **P1/2(10)** (sample taken from Table 3, no. 4).




Fig. S7 ^{13}C NMR spectra of CDCl_3 solutions of (A) **1**, (B) **2(10)**, (C) a mixture of **5–8** and (D) **P1/2(10)** (sample taken from Table 3, no. 4). The solvent peaks were marked with asterisks.




Fig. S8 ^1H NMR spectra of CDCl_3 solutions of (A) 1, (B) 3 and (C) P1/3 (sample taken from Table 3, no. 5).




Fig. S9 ^{13}C NMR spectra of CDCl_3 solutions of (A) 1, (B) 3 and (C) P1/3 (sample taken from Table 3, no. 5). The solvent peaks were marked with asterisks.




Fig. S10 Size distribution of nanoparticles of P1/3 suspended in 99% aqueous mixture.
Concentration: 10 μ M.




Fig. S11 Plot of quantum yield versus the solvent composition of THF/water mixture of P1/3.