

Neutral Linear Amphiphilic Homopolymers Prepared by Atom Transfer Radical Polymerization

Yi Wang^a, Alina O. Alb^b, Jibao He^c, and Scott M. Grayson^{a*}



^a Department of Chemistry, Tulane University, 6400 Freret St., New Orleans, LA, 70118, U.S.A. Fax: 504-865-5596; Tel: 504-862-8135; E-mail: sgrayson@tulane.edu

^b Department of Physics, Tulane University, 6400 Freret St., New Orleans, LA, 70118, U.S.A.


^c Coordinated Instrumentation Facility, 605 Lindy Boggs Building, Tulane University, 6400 Freret St., New Orleans, LA, 70118, U.S.A.

Table of Contents


1) ^1H NMR of amphiphilic monomer 3	S1
2) COSY NMR of amphiphilic monomer 3	S2
3) ^{13}C NMR of amphiphilic monomer 3	S3
4) DEPT NMR of amphiphilic monomer 3	S4
5) ^1H NMR of amphiphilic homopolymer 4	S5
6) ^{13}C NMR of amphiphilic homopolymer 4	S6




S2 COSY NMR (in CDCl_3 , δ) of amphiphilic monomer 3.




DEPT135 NMR Spectrum of monomer in CDCl₃ at 21 °C



S4 DEPT NMR (in CDCl_3 , δ) of amphiphilic monomer 3.



S5 ^1H NMR (in CDCl_3 , δ) of amphiphilic homopolymer 4.



S6 ^{13}C NMR (in CDCl_3 , δ) of amphiphilic homopolymer 4.