

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE

Ethylene/4-methyl-1-pentene copolymers of complex chain architectures by α -diimine Ni(II) catalysts: synthesis, ^{13}C NMR assignment and understanding the chain-walking mechanism

Simona Losio*, Giuseppe Leone, Fabio Bertini, Giovanni Ricci, Maria Carmela Sacchi and Antonella Caterina Boccia*

Received (in XXX, XXX) Xth XXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXX 20XX

DOI: 10.1039/b000000x




Figure S1. ^1H NMR spectrum of E/4M1P copolymer (run 6 in Table 1)

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE




Figure S2. ^1H - ^1H COSY spectrum of E/4MIP copolymer (run 6 in Table 1).

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE




Figure S3. ^1H - ^1H TOCSY spectrum of E/4M1P copolymer (run 6 in Table 1).

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE




Figure S4. T_g plots for runs 1-6




Figure S5. Ethyl branches distribution

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE




Figure S6. Propyl branches distribution




Figure S7. Butyl branches distribution

