

## Supporting Information

### Amphiphilic/Fluorous Random Copolymers as a New Class of Non-Cytotoxic Polymeric Materials for Protein Conjugation

Yuta Koda,<sup>a,b</sup> Takaya Terashima,<sup>a\*</sup> Mitsuo Sawamoto,<sup>a</sup> and Heather D. Maynard<sup>b\*</sup>

\*: corresponding authors

a) Department of Polymer Chemistry, Graduate School of Engineering, Kyoto University  
Katsura, Nishikyo-ku, Kyoto 615-8510, Japan

Tel/Fax: +81-75-383-2603, E-mail: terashima@living.polym.kyoto-u.ac.jp

b) Department of Chemistry and Biochemistry, University of California  
607 Charles E. Young Drive East, Los Angeles, CA 90095-1569, USA


Tel: +1-310-267-5162, E-mail: maynard@chem.ucla.edu


---

#### Contents


<b>Figure S1.</b>	<sup>1</sup> H NMR spectra of <b>P2</b> and (b) <b>P3</b> in acetone- <i>d</i> <sub>6</sub>	S2
<b>Figure S2.</b>	<sup>19</sup> F NMR spectra of <b>P2</b> and (b) <b>P3</b> in acetone- <i>d</i> <sub>6</sub>	S3
<b>Figure S3.</b>	UV-vis spectra of <b>P1</b> after the terminal transformation	S3

## Supporting Data





**Figure S2.**  $^{19}\text{F}$  NMR spectra of (a) **P2** and (b) **P3** in acetone- $d_6$  at 25 °C.



**Figure S3.** UV-Vis spectra of **P1** (black) and **P1** after the transformation of the terminal trithiocarbonate group with AIBN (red):  $[\text{P1}]_0 = 10 \text{ mg/mL}$  in  $\text{H}_2\text{O}$  at 25 °C.