

Supplementary Information for

La(OTf)₃-mediated self-organization of guanosine having alkynyl-Au(I)PPh₃ moiety to induce Au(I)-Au(I) interaction

Xiangtai Meng,^a Toshiyuki Moriuchi,*^a Yuki Sakamoto,^a Masatoshi Kawahata,^b Kentaro Yamaguchi^b and Toshikazu Hirao*^a

^aDepartment of Applied Chemistry, Graduate School of Engineering,
Osaka University,

Yamada-oka, Suita, Osaka 565-0871, Japan

E-mail: moriuchi@chem.eng.osaka-u.ac.jp, hirao@chem.eng.osaka-u.ac.jp

^b Pharmaceutical Sciences at Kagawa Campus, Tokushima Bunri University

1314-1 Shido, Sanuki, Kagawa 769-2193, Japan.

E-mail: yamaguchi@kph.bunri-u.ac.jp




Figure S1. ¹H NMR of GPhAuPPh₃ (CDCl₃, 400 MHz).




Figure S2. ^{13}C NMR of GPhAuPPh_3 (CDCl_3 , 400 MHz).




Figure S3. ^{31}P NMR of GPhAuPPh_3 (CDCl_3 , 109 MHz).




Figure S4. HSQC of GPhAuPPh₃ (CDCl₃, 400 MHz).




Figure S5. HMBC of GPhAuPPh₃ (CDCl₃, 400 MHz).




Figure S6. The CSI-MS spectrum of the octamer of GPhAuPPh₃ with KPF₆.




Figure S7. The CSI-MS spectrum of the octamer of GPhAuPPh_3 with $\text{La}(\text{OTf})_3$.