

SUPPORTING INFORMATION

Partial oxidation of aromatic alcohols via TiO₂ photocatalysis: The influence of substituent groups on the activity and selectivity

Sedat Yurdakal*^a, Vincenzo Augugliaro*^b

Dr. S. Yurdakal, Kimya Bölümü, Fen-Edebiyat Fakültesi, Afyon Kocatepe Üniversitesi, AhmetNecdetSezerKampüsü, 03100, Afyonkarahisar, Turkey, Tel: +90 272 228 1339/230, Fax: +90 0 272 228 12 35, E-mail: sedatyurdakal@gmail.com,

^b "Schiavello-Grillone" Photocatalysis Group, Dipartimento di Ingegneria Elettrica, Elettronica e delle Telecomunicazioni, di tecnologie Chimiche, Automatica e modelli Matematici, University of Palermo, Viale delle Scienze, 90128 Palermo, Italy. Email: vincenzo.augugliaro@unipa.it




Figure S1. SEM images (magnification: x200000) of HPRT and Sigma-Aldrich (SA) photocatalysts, respectively.






Figure S2. TEM images (magnification: X135000 and, X43000 respectively) of HPRT and SA photocatalysts, respectively.



(a)



(b)

Figure S3. XRD patterns of commercial TiO_2 (SA) (a) and HPRT (b) in the presence of CaF_2 (1:1, w/w).

Acknowledgment

The author likes to thank AUBIBAM research centre (Eskisehir, Turkey) for performing TEM analysis.