Supplementary Material (ESI) for **RSC Advances** This journal is © The Royal Society of Chemistry

Supporting Information

L-Proline-catalyzed three-component domino reactions in the synthesis of novel densely functionalized pyrazolo[3,4-b]pyridines

Pethaiah Gunasekaran, Sethuraman Indumathi and Subbu Perumal^a

^aDepartment of Organic Chemistry, School of Chemistry, Madurai Kamaraj University, Madurai 625021, India. Fax: 91-452-2459845. E-mail: subbu.perum@gmail.com.

Supporting Information

- (i) NMR discussion
- (ii) Copies of NMR spectra and ESI-MS for compounds 4a-4w.
- (iii) Copies of ¹⁹FNMR spectra for eight representative compounds.
- (iv) Mechanism study NMR spectra.

NMR discussion

1. Structure determination of 4a.

The structure of 3-methyl-1-aryl-1*H*-pyrazolo[3,4-*b*]pyridine derivatives **4** is in accord with elemental analysis and spectroscopic (1 H, 13 C and 2D NMR) data as illustrated for a representative example **4a**.

$$\begin{array}{c} CI \\ 7.20, d, 1H, J=3.9 \text{ Hz} \\ 134.9 \\ 127.9 \\ 1 \end{array}$$

$$\begin{array}{c} 7.20, d, 1H, J=3.9 \text{ Hz} \\ 134.9 \\ 14.7 \\ 143.3 \\ 143.3 \\ 143.3 \\ 1 \end{array}$$

$$\begin{array}{c} 7.65, d, 1H, J=3.6 \text{ Hz} \\ 135.6 \\ 1 \end{array}$$

$$\begin{array}{c} 7.20, d, 1H, J=3.9 \text{ Hz} \\ 134.9 \\ 144.6 \\ 1 \end{array}$$

$$\begin{array}{c} 1.16.3 \\ 143.3 \\ 143.3 \\ 1 \end{array}$$

Figure 1. Selected 1H and 13C chemical shifts of 4a

In the 1 H NMR spectrum of $\mathbf{4c}$, the H-3" gives a doublet at 7.20 ppm (J=3.9 Hz), which shows (i) a C,H-COSY correlation with carbon signal at 134.9 ppm, assigning it to C-3" and (ii) HMBCs (Figure 2) with C-5a, C-2", C-4" at 185.4, 144.6 and 127.9 ppm respectively. Similarly, the H-4" gives a triplet at 7.01 ppm (J=3.9 Hz), which shows (i) a C,H-COSY correlation with carbon signal at 127.9 ppm assigning it to C-4" and (ii) HMBCs with C-2" and C-5" at 144.6 and 135.6 ppm respectively. The H-5" gives a doublet at 7.65 ppm (J=3.6 Hz) and C,H-COSY correlation with carbon signal at 135.6 ppm, besides showing HMBCs with C-2" at 144.6 and 127.9 ppm respectively. The methyl hydrogens of the pyrazole sub-structure appear as a singlet at 2.12 ppm and show HMBCs with the signals at 143.4 and 116.3 ppm assignable to C-3 and C-3b respectively. The mass spectrum of $\mathbf{4c}$ has a \mathbf{M}^{+} peak at 498.1 in accord with the proposed structure. A single crystal X-ray crystallographic study of $\mathbf{4o}$ confirmed the structure deduced from NMR spectroscopic data (Figure 1).

2. Structural determination of 5r

Methoxy protons of **5r** appear as a singlet at 3.83 ppm in ¹H-NMR and the corresponding carbon appears at 55.4 ppm. This carbon signal shows a HMBC with C-4''. The pyrazole methyl hydrogens appear as a singlet at 1.99 ppm and show HMBCs with the signals at 146.1 and 105.6 ppm assignable to C-3 and C-3b respectively. H-4 appears as a triplet at 4.19 ppm, this multiplicity arises from coupling with the two diastereotpic 5-CH₂ hydrogens which occur as a multiplet. By a C,H-COSY correlation with the signal at 4.19 ppm, the C-4 is assigned to signal at 34.6 ppm. H-4 gives the HMBCs with C-3, C-3b and C-6 at 146,1, 105.6 and 165.1 respectively. Other HMBC correlations shown in Figure 2 support the structure **5r**.




Figure 2. Selected ¹H and ¹³C chemical shifts of **5r**

3. Structural determination of 4x.

The pyrazole methyl hydrogens appear as a singlet at 2.13 ppm, which shows HMBCs with the signals at 143.4 and 116.3 ppm assignable to C-3 and C-3b respectively. Methoxy protons of anisole appear as a singlet at 3.77 ppm and the corresponding carbon occurs at 55.1 ppm, assigned by C,H-COSY spectrum. The signals assigned and the HMBC correlations support the structure and stereochemistry of 4x.









Figure 3. Selected ¹H and ¹³C chemical shifts of **4x**























Figure 16. H,H-COSY spectrum of 4a















Figure 21. C,H-COSY spectrum of 4c-expansion

















































































































Figure 90. ¹³C NMR spectrum of 5r-expansion









Figure 91. Dept-135 NMR spectrum of 5r



































Figure 113.DEPT-135 spectrum of 4x










Figure 116. C,H-COSY spectrum of 4x




Figure 119. HMBC spectrum of 4x




Figure 120. HMBC spectrum of 4x-expansion1








Figure 121. HMBC spectrum of 4x -expansion2









Mechanism study NMR spectrums -intermediate identification from the reaction





Figure 126. reaction mixture-1



Electronic Supplementary Material (ESI) for RSC Advances This journal is © The Royal Society of Chemistry 2013