

Convenient synthesis of C75, an inhibitor of FAS and CPT1

Carolina Sánchez,^a Kamil Makowski,^{a,b} Paula Mera,^b Jaume Farràs,^a Ernesto Nicolás,^a Laura Herrero,^b Guillermina Asins,^b Dolors Serra,^b Fausto G. Hegardt,^b Xavier Ariza,^{*a} and Jordi Garcia^{*a}

^aDepartament de Química Orgànica, CIBEROBn and IBUB, Facultat de Química, Universitat de Barcelona, C/Martí i Franquès 1-11, 08028-Barcelona (Spain).

^bDepartament de Bioquímica i Biologia Molecular, CIBEROBn and IBUB, Facultat de Farmàcia, Universitat de Barcelona, C/Joan XXIII s/n, 08028-Barcelona (Spain)

E-mail: xariza@ub.edu, jordigarciagomez@ub.edu

Supporting Information Index

¹ H NMR and ¹³ C NMR spectra of 1	2
¹ H NMR and ¹³ C NMR spectra of 2	3
¹ H NMR and ¹³ C NMR spectra of <i>syn</i> - 5	4
¹ H NMR and ¹³ C NMR spectra of <i>trans</i> - 8	5
¹ H NMR and ¹³ C NMR spectra of <i>cis</i> - 8	6
¹ H NMR and ¹³ C NMR spectra of <i>trans</i> - 9	7
¹ H NMR and ¹³ C NMR spectra of <i>cis</i> - 9	8
¹ H NMR and ¹³ C NMR spectra of <i>trans</i> - 4	9
¹ H NMR and ¹³ C NMR spectra of C75	10
¹ H NMR and ¹³ C NMR spectra of 3	11
¹ H NMR and ¹³ C NMR spectra of <i>cis,trans</i> -2-octyl-5-oxo-4-[(phenylselanyl)methyl]tetrahydrofuran-3-carboxylic acid	12





















