

Supporting information

Synthesis of branched-phosphodiester and mannose-centered fucosylated glycoclusters and their binding studies with *Burkholderia ambifaria* lectin (BambL)

Caroline Ligeour,^[a] Aymeric Audfray,^[b] Emilie Gillon,^[b] Albert, Meyer,^[a] Sébastien Vidal,^[c] Jean-Jacques Vasseur,^[a] Anne Imberty,^{*[b]} François Morvan^{*[a]}

[a] Institut des Biomolécules Max Mousseron UMR 5246 CNRS Université Montpellier 1 Université Montpellier 2, Université Montpellier 2, CC1704, Place E. Bataillon, 34095 Montpellier Cedex 5(France)

[b] CERMAV - CNRS (affiliated with Université Joseph Fourier and ICMG), BP 53, 38041, Grenoble (France)

[c] Institut de Chimie et Biochimie Moléculaires et Supramoléculaires, Laboratoire de Chimie Organique 2 - Glycochimie, UMR 5246, Université Lyon 1 and CNRS, 43 Boulevard du 11 Novembre 1918, 69622, Villeurbanne (France)

morvan@univ-montp2.fr

anne.imberty@cermav.cnrs.fr

Table of contents

Figure S1. SPR Sensorgrams for compounds 1, 2, 4 and 5.....	S3
Figure S2. ITC binding isotherm of 1	S4
Figure S3. ITC binding isotherm of 2	S4
Figure S4. ITC binding isotherm of 3	S5
Figure S5. ITC binding isotherm of 4	S5
Figure S6. ITC binding isotherm of 5	S6
Figure S7. ^1H NMR of compound 9a	S7
Figure S8. ^{13}C NMR of compound 9a	S8
Figure S9. ^1H NMR of compound 9b	S9
Figure S10. ^{13}C NMR of compound 9b	S10
Figure S11. ^1H NMR of compound 9c	S11
Figure S12. ^{13}C NMR of compound 9c	S12
Figure S13. ^1H NMR of compound 14	S13
Figure S14. ^{13}C NMR of compound 14	S14
Figure S15. ^1H NMR of compound 15	S15
Figure S16. ^{13}C NMR of compound 15	S16
Figure S17. ^1H NMR of compound 1	S17
Figure S18. ^{13}C NMR of compound 1	S18
Figure S19. ^1H NMR of compound 2	S19
Figure S20. ^{13}C NMR of compound 2	S20
Figure S21. ^1H NMR of compound 3	S21
Figure S22. ^{13}C NMR of compound 3	S22

Figure S23. ^1H NMR of compound 4 S23

Figure S24. ^{13}C NMR of compound 4 S24

Figure S25. ^1H NMR of compound 5 S25

Figure S26. ^{13}C NMR of compound 5 S26






Figure S1. SPR Sensorgrams for compounds 1, 2, 4 and 5




$[1] = 0.15 \text{ mM}$, $[\text{BambL}] = 0.0125 \text{ mM}$

Figure S2. ITC binding isotherm of **1**




$[2] = 0.15 \text{ mM}$, $[\text{BambL}] = 0.0125 \text{ mM}$

Figure S3. ITC binding isotherm of **2**




[3] = 0.075 mM, [BambL] = 0.0130 mM

Figure S4. ITC binding isotherm of 3



[4] = 0.15 mM, [BambL] = 0.0133 mM

Figure S5. ITC binding isotherm of 4



[**5**] = 0.075 mM, [BambL] = 0.0130 mM

Figure S6. ITC binding isotherm of 5




Figure S7. ¹H NMR of compound 9a




Figure S8. ^{13}C NMR of compound 9a




Figure S9. ^1H NMR of compound 9b




Figure S10. ^{13}C NMR of compound 9b




Figure S11. ¹H NMR of compound 9c




Figure S12. ^{13}C NMR of compound 9c




Figure S13. ^1H NMR of compound 14




Figure S14. ^{13}C NMR of compound 14




Figure S15. ¹H NMR of compound 15




Figure S16. ^{13}C NMR of compound 15




Figure S17. ^1H NMR of compound 1




Figure S18. ^{13}C NMR of compound 1




Figure S19. ¹H NMR of compound 2




Figure S20. ^{13}C NMR of compound 2




Figure S21. ^1H NMR of compound 3




Figure S22. ^{13}C NMR of compound 3




Figure S23. ^1H NMR of compound 4




Figure S24. ^{13}C NMR of compound 4




Figure S25. ^1H NMR of compound 5




Figure S26. ^{13}C NMR of compound 5