

# Time-Dependent Density Functional Study of UV-Visible Absorption Spectra of Small Noble Metal Clusters ( $\text{Cu}_n$ , $\text{Ag}_n$ , $\text{Au}_n$ , $n = 2 - 9, 20$ )

Berkahem Anak<sup>a</sup>, Mustapha Bencharif<sup>a</sup>, Franck Rabilloud<sup>b</sup>

<sup>a</sup> Laboratoire de Chimie des Matériaux, Université Constantine 1,  
Constantine, Algérie

<sup>b</sup> Institut Lumière Matière, UMR5306 Université Lyon 1 - CNRS, Université de Lyon,  
69622 Villeurbanne Cedex, France  
franck.rabilloud@univ-lyon1.fr

November 21, 2013




Figure 7: Absorption spectra of gold and copper clusters.




Figure 8: Absorption spectra of  $\text{Ag}_4$  ( $D_{2h}$ ),  $\text{Ag}_6$  ( $D_{3h}$ ) and  $\text{Ag}_8$  ( $T_d$ ) clusters using the SAOP method with or without the spin-orbit (SO) coupling.




Figure 9: Absorption spectra of  $\text{Cu}_4$  ( $D_{2h}$ ),  $\text{Cu}_6$  ( $D_{3h}$ ) and  $\text{Cu}_8$  ( $T_d$ ) clusters using the SAOP method with or without the spin-orbit (SO) coupling.