

Supporting Information for

Degradable, Silyl Ether, Thiol-Ene Networks

Taylor Ware ^{††1}, Abby R. Jennings ^{‡2}, Zahra S. Bassampour ², Dustin Simon ¹, David Y. Son ^{*2}
and Walter Voit ^{*1}

¹ *Department of Materials Science and Engineering, The University of Texas at Dallas, 800 West
Campbell RD, RL 3 Richardson, TX, 75080, USA*

² *Department of Chemistry, Center for Drug Discovery, Design, and Delivery (CD4), Southern
Methodist University, Dallas, TX, 75275, USA*

Corresponding Authors*: dson@mail.smu.edu, walter.voit@utdallas.edu

Contents	Figure
1. ¹ H, ¹³ C, and ²⁹ Si NMR spectra for 6 and 7	S1-S6
2. Mass Spectra for 9 , the alpha product, 3 , and 11	S7-S10




Figure S 1. ^1H NMR (500 MHz, CDCl_3) compound **6**.




Figure S 2. ^{13}C NMR (125 MHz, CDCl_3) compound **6**.




Figure S 3. ^{29}Si -NMR (99 MHz, CDCl_3) compound **6**.




Figure S 4. ¹H NMR (500 MHz, CDCl₃) compound 7.




Figure S 5. ¹³C NMR (125 MHz, CDCl₃) compound 7.




Figure S 6. ²⁹Si-NMR (99 MHz, CDCl₃) compound 7.




Figure S 7. Mass spectrum of compound **9**.




Figure S 8. Mass spectrum of the alpha product.




Figure S 9. Mass spectrum of compound **3**.




Figure S 10. Mass spectrum of compound **11**.