

Supporting Information for

Rapid assembly of quinazolinone scaffold via copper-catalyzed tandem reaction of 2-bromobenzamides with aldehydes and aqueous ammonia: application to the synthesis of alkaloid tryptanthrin

Shanghai Guo,*^a Yan Li,^b Li Tao,^a Wenwen Zhang^a and Xuesen Fan*^a


^a Collaborative Innovation Center of Henan Province for Green Manufacturing of Fine Chemicals, Key Laboratory of Green Chemical Media and Reactions, Ministry of Education, School of Chemistry and Chemical Engineering, Henan Normal University, Xinxiang, Henan 453007, P. R. China. E-mail: shguo@htu.cn; xuesen.fan@htu.cn


^b School of Science, Jiaozuo Teachers' College, Jiaozuo, Henan 450001, P. R. China


Table of Content


I. Copies of ^1H and ^{13}C NMR spectra of 3a-3w	P2-P24
II. Copies of ^1H and ^{13}C NMR spectra of 5a-5l	P25-P36
III. Copies of ^1H and ^{13}C NMR spectra of 6 and 7	P37-P38
IV. Copies of ^1H and ^{13}C NMR spectra of 9	P39


I. Copies of ^1H and ^{13}C NMR spectra of 3a-3w






































































II. Copies of ^1H and ^{13}C NMR spectra of 5a-5l



































III. Copies of ^1H and ^{13}C NMR spectra of 6 and 7





IV. Copies of ^1H and ^{13}C NMR spectra of 9

