

Supplemental figure

Simultaneous determination of aconite alkaloids and ginsenosides using online solid phase extraction hyphenated with polarity switching UHPLC–MS–MS

Yuelin Song ^{1#}, Na Zhang ^{1,2,3#}, Yong Jiang ⁴, Jun Li ¹, Yunfang Zhao ¹, Shepo Shi ^{1*}, Pengfei Tu ^{1,4*}

¹. *Modern Research Center for Traditional Chinese Medicine, Beijing University of Chinese Medicine, Beijing 100029, China*

². *School of Chinese Materia Medica, Beijing University of Chinese Medicine, Beijing 100102, China*

³. *Baotou Medical College, Baotou 014060, China*

⁴. *State Key Laboratory of Natural and Biomimetic Drugs, School of Pharmaceutical Sciences, Peking University, Beijing 100191, China*

#: These two authors contribute equally to this article.

*Corresponding authors.

Tel./fax: +86-10-64286350; *E-mail address*: shishepo@163.com (S.P. Shi).

Tel./fax: +86-10-82802750; *E-mail address*: pengfeitu@163.com (P.F. Tu).

Figure legend:

Fig. S1 MS² spectra of all the reference compounds under positive and negative modes for aconite alkaloids and ginsenosides, respectively. For ginsenosides, the formate anions were selected as the precursor ions, while the positive quasi-molecular ions were set as the parent ions for alkaloids.





