

Supporting Information


Imidazolium Salt-supported Mukaiyama Reagent: An Efficient Condensation Reagent for Amide Bond Formation



Khima Pandey, Sunita Choudhary, Manoj Kumar Muthyala and Anil Kumar*



Department of Chemistry, Birla Institute of Technology and Science, Pilani, 333031, India

E-mail: anilkumar@pilani.bits-pilani.ac.in



Copies of ^1H NMR and ^{13}C NMR spectra of **13**, **14**, **15**, **18a-p**, **20**











7.91
7.89
7.84
7.68
7.66
7.60
7.59
7.57
7.54
7.52
7.50
7.43
7.41
7.39
7.20
7.19
7.17







165.75



137.92
135.02
131.89
129.14
128.84
127.03
124.62
120.20















8.34
8.31
8.04
8.01
7.82
7.53
7.50
7.21
7.18



—2.36




163.45
149.71
140.63
135.15
134.63
129.76
128.23
124.01
120.48

—20.95







8.18
8.15
7.87
7.84
7.31
7.27
7.26
7.23
- 6.62
4.57
4.55




18h







165.37
149.61
139.91
137.46
128.93
128.21
127.96
127.94
123.82
44.45




18h
















165.66, 153.32, 141.21, 137.93, 130.47, 129.10, 124.63, 120.31, 107.92, 104.54, 60.96, 56.37











9.06
8.91
8.67
8.18
8.16
7.64
7.62
7.35
7.33
7.31
7.18
7.16
7.14



164.29
152.11
147.99
137.65
135.61
130.99
129.09
125.05
123.72
120.78

