### **Electronic Supplementary Information**

## Enhanced in vitro cytotoxicity and cellular uptake of DNA bases functionalized gold

### nanoparticles in HeLa cell lines

Shubhangi Borse, Satyawati Joshi\* and Ayesha Khan

Department of Chemistry, Savitribai Phule Pune University, Ganeshkhind Road, Pune 411007

Maharashtra, India

e-mail: ssjoshi@chem.unipune.ac.in

I. Zeta potential data

II. FTIR data

III. Structure of adenine, thymine and cytosine and their pair: Binding interaction with

gold nanoparticles.

IV. Cytotoxicity study of healthy cells line

# I. Zeta-potential data



ESI – Figure 1. Zeta potential of functionalized AuNPs.

Name	Diameter	Zeta	pН
	(TEM)	Potential,	
	nm	mV	
AuNPs	$6.3 \pm 0.04$	-53.2	8.28
AuNPs-A	$4.8 \pm 0.11$	-28.0	7.81
AuNPs-T	$15.5 \pm 0.2$	-41.8	8.32
	3		
AuNPs-C	$7.1 \pm 2.09$	-31.8	8.31
AuNPs-AC	$6.7 \pm 0.14$	-34.0	8.18
AuNPs-AT	$6.1 \pm 0.25$	-37.1	8.10
AuNPs-CT	$9.4 \pm 0.22$	-24.5	8.30

**ESI– Table 1.** Particle diameter (TEM) and zeta potential values and pH of functionalized AuNPs.

#### **III. FTIR data**

The band assignments are reported in this section where the band intensity is indicated as s = strong, m = moderate, w= weak and vw = very weak



ESI – Figure 2. FTIR spectra of cytosine (black line) and AuNPs-Cytosine (red line).

Modes	Free (cm <sup>-1</sup> )	Bound (cm <sup>-1</sup> )
$\nu NH_2$	3202	3360 (s)
v C(2)=O(8)	1703	1645 (s)

v N(3) = C(4)	1536	1504 (m)
v C(5) = C(6)	1460	1437 (m)
$\nu$ C(4)-NH <sub>2</sub>	1370	1378 (s)
vC-NH <sub>2</sub>	960	926 (m)
vN-C-N	779	787 (w)
Ring breathing	696	696 (w)
of Imidazole		

ESI - Table 2. Main vibrational modes of free and bound cytosine



ESI – Figure 3. FTIR spectra of thymine (black line) and AuNPs-thymine (red line).

Modes	Free (cm <sup>-1</sup> )	Bound (cm <sup>-1</sup> )
v N(3)H	3110	3369(s)
ν N(1) H		
v C(2)=O(7)	1720	1753 (s)
v C(2)=O(7)	1653	1670 (s)
δ (NH)	1441	1420 (s)
		1396 (s)
δ (C-O)	1195	1130 (m)
	1030	1037(m)
	933	950 (m)
$\delta$ (C-H(ar) oop)	746	763 (vw)

**ESI – Table 3.** Main vibrational modes of free and bound thymine.



ESI-Figure 4. FTIR spectra of adenine (black line) and AuNPs-adenine (red line).

Modes	Free (cm <sup>-1</sup> )	Bound (cm <sup>-1</sup> )
v NH <sub>2</sub>	3236	3385 (s)
v C-N(7)	1720	1661 (s)
v N-H	1595	1579 (s)
v C-C	1321	1345 (s)
ν C-Η		
v C-N	1238	1245 (s)
ν С-Н		
v C-C-N	1007	1022 (m)
v C-H (In plane)	1103	1092 (vw)
v C-H (Out plane)		
vNH <sub>2</sub>	936	921 (vw)
vN-C-N	1166	
	936	
	896	

ESI – Table 4. Main vibrational modes of free and bound adenine.



**ESI – Figure 5.** FTIR spectra of pair of DNA bases functionalized AuNPs.



**ESI–** Figure 6. DNA bases (cytosine, thymine and adenine) binding interaction with gold nanoparticles. The numbers in the blue color are the positions of the atoms of the bases. Binding interactions are indicated by the blue lines.

7

![](_page_7_Figure_0.jpeg)

IV. Cytotoxicity study of healthy cells line

**ESI–Figure 7.** Cytotoxicity measurements of functionalized AuNPs in healthy cells at various concentrations  $2-30 \mu g/mL$  for 24 h using MTT cell viability assay.

![](_page_8_Figure_0.jpeg)

**ESI–Figure 8.** Cytotoxicity measurements of functionalized AuNPs in healthy cells at various concentrations  $2-30 \ \mu g/mL$  for 48 h using MTT cell viability assay.