

Supporting Information for:

RbFe²⁺Fe³⁺F₆: Synthesis, Structure, and Characterization of a New Charge-Ordered Magnetically Frustrated Pyrochlore-Related Mixed-Metal Fluoride

Sun Woo Kim,¹ Sang-Hwan Kim,¹ P. Shiv Halasyamani^{1,}, Mark Green,² Kanwal Preet Bhatti³, C.
Leighton^{3,*} Hena Das⁴, and Craig Fennie^{4,*}*

¹Department of Chemistry, University of Houston, 136 Fleming Building, Houston, TX 77204-5003.

²Department of Materials Science and Engineering, University of Maryland, College Park, MD, 20742-2115
and NIST Center for Neutron Research, National Institute of Standards and Technology, 100 Bureau Drive,
Gaithersburg, MD 20899-6103.


³Department of Chemical Engineering and Materials Science, University of Minnesota, 151 Amundson Hall,
421 Washington Ave SE, Minneapolis, MN 55455.

⁴ School of Applied & Engineering Physics, Cornell University, Ithaca, NY 14853.


¹psh@uh.edu , ³leighton@umn.edu, ⁴fennie@cornell.edu

- S1. Experimental and calculated powder X-ray diffraction patterns for RbFe₂F₆
- S2. ORTEP (50% probability ellipsoids) diagrams for RbFe₂F₆
- S3. IR spectrum for RbFe₂F₆
- S4. UV-Vis diffuse reflectance spectrum for RbFe₂F₆
- S5. Thermogravimetric analysis and differential thermal analysis diagram for RbFe₂F₆
- S6. Powder X-ray diffraction data for final residuals after TGA Experiment


S1. Experimental and calculated powder X-ray diffraction patterns for RbFe₂F₆



S2. ORTEP (50% probability ellipsoids) diagrams for RbFe₂F₆




S3. IR spectrum for RbFe_2F_6




RbFe_2F_6	
$\nu(\text{Fe-F})$	$\nu(\text{Fe-F-Fe})$
731	408
	463
	503

S4. UV-Vis diffuse reflectance spectrum for RbFe₂F₆



S5. Thermogravimetric analysis and differential thermal analysis diagram for RbFe_2F_6



S6. Powder X-ray diffraction data for final residuals after TGA Experiment.

