

Supplementary Information

**A New Approach to Light up DNA/Ag Nanocluster-based
Beacons for Bioanalysis**

*Libing Zhang, Jinbo Zhu, Zhixue Zhou, Shaojun Guo, Jing Li, Shaojun Dong**
*and Erkang Wang**

State Key Laboratory of Electroanalytical Chemistry, Changchun Institute of Applied
Chemistry, Chinese Academy of Sciences, Changchun, Jilin, 130022, P. R. China, and
University of the Chinese Academy of Sciences, Beijing, 100039, P. R. China.

* Corresponding author: dongsj@ciac.jl.cn; ekwang@ciac.jl.cn

Supplementary Table and Figures

Table S1. Names and sequences of the oligonucleotides.

Oligonucleotide	Sequence
Ag-DNA	5'-CCCTTAATCCCAGCACATCTGATAGTTCTATGCT-3'
En-DNA	5'-GAACTATCAGATGTGCTGGGTGGGGTGGGGTGGGG-3'
B-DNA	5'-TATATCAGCATAGAACTATCAGATGTGCT-3'
T1: T-DNA	5'-AGCACATCTGATAGTTCTATGCTGATATA-3'
T2	5'-AGCACATCTGATAGTTCTATGCTTATATA-3'
T3	5'-AGCACATCTGATTGTTCTATGCTTATATA-3'
T4	5'-AGCATATCTGTTAGTTATATGCTTATATA-3'
B-DNA (Amplification)	5'-TATATCAGCATAGAACTATCAGATGTGCTATGA-3'
T-DNA (Amplification)	5'-TCATAGCACATCTGATAGTTCTATGCTGATATAGTGA-3'
A-Ag-DNA	5'-CCCTTAATCCCAGCACATCTGATAACCACACCAA-3'
A-En-DNA	5'-GTGGTTATCAGATGTGCTGGGTGGGGTGGGGTGGGG-3'
A-B-DNA	5'-ATCAGATGTGCT-3'
Aptamer	5'-GGTTGGTGTGGTTGG-3'

Table S2. The lifetimes of dark DNA/Ag NCs and bright DNA/Ag NCs.

sample	τ_1 (ns)	A_1 (%)	τ_2 (ns)	A_2 (%)	$\tau_{ave}^{[a]}$ (ns)	χ^2
Bright Ag NCs	3.72	97.66	9.20	2.34	3.85	1.100
Dark Ag NCs	3.14	93.65	9.44	6.35	3.54	1.024

[a] The average lifetime was calculated according to $\tau_{ave} = \sum A_i \tau_i$.




Figure S1 The fluorescence intensity of bright DNA/Ag NCs enhanced as a function of incubation time.




Figure S2 TEM image of DNA/Ag NCs.




Figure S3 Native polyacrylamide gel electrophoresis analysis of the behavior of DNA strands with and without clusters. Lane 1: En-DNA; lane 2: Ag-DNA; lane 3: Ag-DNA with clusters; lane 4: En-DNA+Ag-DNA; lane 5: En-DNA+Ag-DNA with clusters.




Figure S4 (A) Fluorescence emission spectra of analyzing different concentrations of T-DNA using the ExoIII-based amplification method (from **a** to **k**): 0, 0.1, 0.5, 1, 2, 5, 10, 20, 50, 100, 200 nM. The concentration of other DNA used is all 1.0 μ M. (B) The relationship between the fluorescence intensity and the concentration of T-DNA. The inset shows a linear relationship ($R = 0.993$) over the concentration of T-DNA from 0.1 to 5.0 nM.




Figure S5 Native 15% polyacrylamide gel analysis of strand displacement mechanism. Different DNA were added into lanes 1-9. Lane 1: A-Ag-DNA; lane 2: A-En-DNA; lane 3: A-B-DNA; lane 4: Aptamer; lane 5: A-Ag-DNA+A-En-DNA; lane 6: A-Ag-DNA+A-B-DNA+Aptamer; lane 7: A-Ag-DNA+A-En-DNA+A-B-DNA+Aptamer; lane 8: A-Ag-DNA+A-En-DNA+A-B-DNA+Aptamer+thrombin; lane 9: Aptamer+thrombin. Concentrations for each DNA in PAGE are all 1.0 μ M.




Figure S6 Fluorescent intensity at 620 nm with different amounts of Exo III; curve a : 0 nM target DNA present, curve b : 100 nM target DNA present.