

Supporting Information for the Manuscript Entitled

Self-assembly of Biodegradable Copolyester and Reactive HPMA-based Polymers into Nanoparticles as an Alternative Stealth Drug Delivery System

Eliézer Jäger,^{†,*} Alessandro Jäger,[†] Tomáš Etrych,[†] Fernando Carlos Giacomelli,[‡]
Petr Chytil,[†] Alexander Jigounov,[†] Jean-Luc Putaux,[§] Blanka Říhová,[‡]
Karel Ulbrich,[†] and Petr Štěpánek^{†,*}

[†] Institute of Macromolecular Chemistry v.v.i., Academy of Sciences of the Czech Republic,
Heyrovsky Sq. 2, 162 06 Prague 6, Czech Republic.

[‡] Centro de Ciências Naturais e Humanas, Universidade Federal do ABC, 09210-170 Santo André,
Brazil.

[§] Centre de Recherches sur les Macromolécules Végétales (CERMAV-CNRS), BP 53, 38041 Grenoble
Cedex 9, France - affiliated with Joseph Fourier University and member of the Institut de Chimie
Moléculaire de Grenoble.

[‡] Institute of Microbiology v.v.i., Academy of Sciences of the Czech Republic, Vídenska 1083, 142 20
Prague 4, Czech Republic.

Corresponding Authors: jager@imc.cas.cz Tel: +420 296 809 322
stepanek@imc.cas.cz Tel: +420 296 809 211




Figure S1. SEC chromatograms for PBS/PBDL (a) and HPMA-chol (b).




Figure S2. ¹H and ¹³C NMR of the poly(butylene succinate-co-butylene dilinoleate) aliphatic copolyester.




Figure S3. PBS/PBDL ¹³C NMR spectra showing the divide of carbonyl groups (C=O) around 64 ppm.




Figure S4. Distribution of R_H as revealed by REPES algorithm from the autocorrelation function $g_2(t)$ measured at 90° for NP1:2.5 in water.