

Supporting Information for:

New organic dyes based on dibenzofulvene bridge for highly efficient dyes–sensitized solar cells.

Agostina Lina Capodilupo,^a Luisa De Marco,^c Eduardo Fabiano,^a Roberto Giannuzzi,^c Giuseppina Anna Corrente,^b Angela Scrascia,^a Claudia Clarlucci,^a Maria Pia Cipolla,^c Giuseppe Gigli^{acd} and Giuseppe Ciccarella.^{ad}

^a Istituto Nanoscienze – CNR, National Nanotechnology Laboratory (NNL), Via Arnesano, 73100 Lecce, Italy. E-mail: giuseppe.ciccarella@unisalento.it

^b Università della Calabria, via Pietro Bucci, 87036 Arcavacata di Renze, Cosenza, Italy

^c Center for Biomolecular Nanotechnologies (CBN) Fondazione Istituto Italiano di Tecnologia (IIT), Via Barsanti 1, Arnesano, 73010, Italy

^d Dipartimento di Matematica e Fisica “Ennio De Giorgi”, Università del Salento, Via Monteroni, 73100, Lecce, Italy.

^e Dipartimento di Ingegneria dell’Innovazione, Università del Salento, Via Monteroni, 73100, Lecce, Italy.








Figure S1 Cyclic voltammetry of the dyes.








2








TK1













4








TK2








6





TK3

