

Bisphosphonate–polyaspartamide conjugates as bone targeted drug delivery systems

D. Paolino,^a M. Licciardi,^b C. Celia,^c G. Giammona,^b M. Fresta^a and G. Cavallaro^{*b}

^aDepartment of Health Sciences, IRC FSH-Interregional Research Center for Food Safety & Health, University of Catanzaro “Magna Graecia”, Campus Universitario “S. Venuta”, Viale S. Venuta, Germaneto, I-88100 Catanzaro, Italy.

^bDepartment of Biological, Chemical and Pharmaceutical Sciences and Technologies, (STEBICEF), Sezione di Chimica e Tecnologie Farmaceutiche, Università degli Studi di Palermo, Via Archirafi 32, 90123 Palermo, Italy.

^cDepartment of Pharmacy, University of Chieti – Pescara “G. d’Annunzio”, Via dei Vestini 31, Chieti 66013, Italy, and Department of Nanomedicine, The Houston Methodist Research Institute, 6670 Bertner Avenue, Houston, TX, 77030, USA.

Table S1. The average weight of bone tissues and organs collected from rats and processed according to Materials and Methods section.

Samples	Average weight (g)
Blood	2.16
Heart	1.3
Lung	1.6
Liver	11.0
Spleen	0.78
Kidneys	2.26
Sternum	2
Femur	1.2
Skeleton	90