

Supporting information

Bioreducible Dextran-Polyethylenimine Conjugates Regulate Transgene Expression Distribution In Vivo

Jian Zhao^{1*}, Chunxiao Wang^{1,2}, Peng Zhao², Xuejun Wen², and Chao Lin^{2*}

¹Key Laboratory of Rubber-Plastics Ministry of Education and Shandong Provincial
Key Laboratory of Rubber-Plastics, Qingdao University of Science & Technology,
Qingdao, 266042, People's Republic of China

²Shanghai East Hospital, The Institute for Biomedical Engineering and Nanoscience,
Tongji University School of Medicine, Tongji University, Shanghai, 200092, People's
Republic of China.

Fax: 0086 21 65983706-0; Tel: 0086 21 65988029.

* Corresponding authors.

E-mail: chaolin@tongji.edu.cn (CL),

zp@tongji.edu.cn (ZP),

and zhaojian@gmail.com(JZ)




Figure S1. a) GPC profile of Dex5k-SS-LPEI2k and 2kDa LPEI disulfide pyridine; b) GPC profile of Dex5k-SS-LPEI2k in the presence of 2.5 mM dithiothreitol (DTT)




Figure S2. FT-IR spectrum of dextran, 2kDa PEI disulfide pyridine and Dex10k-SS-PEI5k.