

Supporting Information for

**Design of Multi-functional Linear Polymers that Capture and  
Neutralize a Toxic Peptide: A Comparison with Cross-linked  
Nanoparticles**

*Y. Wada<sup>a</sup>, H. Lee<sup>a</sup>, Y. Hoshino<sup>a\*</sup>, S. Kotani<sup>b</sup>, K. J. Shea<sup>c</sup>, and Y. Miura<sup>a</sup>*


<sup>a</sup> Department of Chemical Engineering, Kyushu University, 744 Motoooka, Nishi-ku, Fukuoka  
819-0395, Japan

<sup>b</sup> Priority Organization for Innovation and Excellence, Kumamoto University, Kumamoto  
862-0973, Japan

<sup>c</sup> Department of Chemistry, University of California, Irvine, Irvine, CA 92697, USA

\*To whom all correspondence should be addressed. Phone: +81-92-2769. Fax: +81-92-2769. E-mail:

yhoshino@chem-eng.kyushu-u.ac.jp



**Figure S1.** Binding isotherms of LP<sub>1000-11</sub> (a), LP<sub>300-5</sub> (b), LP<sub>300-10</sub> (c) and LP<sub>300-11</sub> (d) to Melittin immobilized QCM in PBS at 37 °C.