Supporting Information

Thiophenoazomethines: Electrochromic Materials Exhibiting Visible-to-Near-IR Color Changes and Tuneable Colors Contingent on Degree of Conjugation

Satyananda Barik,[1,2†] Daminda Navarathne,[1†] Maxence LeBorgne,[1,3] W. G. Skene[1*]

[1]Laboratoire de caractérisation photophysique des matériaux conjugués
Département de Chimie, Pavillon JA Bombardier,
Université de Montréal, CP 6128, succ. Centre-ville,
Montréal, Québec, Canada H3C 3J7

[2]Current address: Institute of Chemical and Engineering Sciences
1 Pesek Road, Jurong Island
Singapore-627833
Singapore

[3]Current address: Département de chimie
Institut Universitaire de Technologie Besançon-Vesoul
30 Avenue de l'Observatoire - BP 1559 – 25009
Besançon cedex France

[†]Contributed equally to the manuscript

Corresponding Author Email: w.skene@umontreal.ca

Table of Contents

Figure S1. Absorbance spectra of thin-film of **7 (——)**, **8** (——), **9 (——)**, **10** (——) and **11**(——) on glass substrates.

Figure S2. Absorbance spectra of neutral **8** (━) in solution recorded with the addition of $FeCl_3$ (━) followed by the addition of hydrazine hydrate (━).

Figure S3. Absorbance spectra of **8** (━) with the addition of trifluoroacetic acid (━) followed by the addition of triethylamine (━).

Figure S4. Gel permeation elugrams of **7** (■), **8** (●), **9** (▲), **10** (▼) and **11** (◆) measured in THF.

Figure S5. Anodic cyclic voltammograms of **9** measured in dichloromethane with TBAPF$_6$ as the supporting electrolyte.

Figure S6. Anodic cyclic voltammograms of **10** measured in dichloromethane with TBAPF$_6$ as the supporting electrolyte.

Figure S7. Cyclic voltammogram (—) and differential pulse voltammograms (—) of **11** in 10^{-4}M polymer solution in 0.1 M TBAPF$_6$/dichloromethane with ferrocene as the internal standard. The potentials are reported against Fc/Fc$^+$.

Figure S8. ^{1}H NMR spectrum of **1** recorded in acetone-d$_6$.

Figure S9. ^{1}H NMR spectrum of **3** recorded in CDCl$_3$.

Figure S10. ^{1}H NMR spectrum of **6** recorded in CDCl$_3$.

Figure S11. ^{1}H NMR spectrum of **9** recorded in CDCl$_3$.

Figure S12. ^{1}H NMR spectrum of **10** measured in CDCl$_3$.

Figure S13. ^{1}H NMR spectrum of **7** recorded in acetone-d$_6$.

Figure S14. ^{13}C NMR spectrum of **7** recorded in acetone-d$_6$.

Figure S15. ^{1}H NMR spectrum of **8** recorded in acetone-d$_6$.

Figure S16. ^{13}C NMR spectrum of **8** recorded in acetone-d$_6$.

Figure S17. ^{1}H NMR spectrum of **11** recorded in acetone-d$_6$.