

Influence of high magnetic field on electric-dipole emission spectra of Eu^{3+} in different single crystals

Guihuan Du, Peng Liu, Weiwei Guo, Yibo Han, Junpei Zhang, Zongwei Ma, Junbo Han, Zuli Liu and Kailun Yao




Fig 1 PL spectra of $^5\text{D}_0\text{-}^7\text{F}_2$ band of $\text{GdVO}_4:\text{Eu}^{3+}$ excited by 396nm laser under different magnetic field at 4.2K.




Fig 2 PL spectra of $^5\text{D}_0\text{-}^7\text{F}_2$ band of $\text{YVO}_4:\text{Eu}^{3+}$ excited by 396nm laser under different magnetic field at 4.2K.




Fig 3 PL spectra of 5D_0 - 7F_4 band of GdVO₄:Eu³⁺ excited by 396nm laser under different magnetic field at 4.2K.




Fig 4 PL spectra of 5D_0 - 7F_4 band of YVO₄:Eu³⁺ excited by 396nm laser under different magnetic field at 4.2K.




Fig 5 Integrated intensity of ${}^5D_0-{}^7F_2$ PL of GdVO₄:Eu³⁺ single crystal versus magnetic field at 4.2K




Fig 6 Integrated intensity of ${}^5D_0-{}^7F_2$ PL of YVO₄:Eu³⁺ single crystal versus magnetic field at 4.2K




Fig 7 PL of 5D_0 - 7F_2 in YVO₄:Eu³⁺ single crystal in different direction for parallel and vertical magnetic field direction at 77K




Fig 8 PL of 5D_0 - 7F_2 in GdVO₄:Eu³⁺ single crystal in different direction for parallel and vertical magnetic field direction at 77K