

Cellular Responses of Aniline Oligomers: A Preliminary Study

Xiaoyong Zhang^{a,#}, Hongxu Qi^{a,#}, Shiqi Wang^a, Lin Feng^a, Yan Ji^a, Lei Tao^{a,*}, Shuxi Li^a,
Yen Wei^{a,b,*}

Received (in XXX, XXX) Xth XXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXX 20XX

DOI: 10.1039/b000000x

^a Department of Chemistry and ^b Key Lab of Organic Optoelectronic & Molecular Engineering of Ministry of Education, Tsinghua University, Beijing, 100084, P. R. China.
leitao@mail.tsinghua.edu.cn; weiyen@tsinghua.edu.cn

Results and Discussion




Fig. S1 Mass spectrum reports of (A) aniline trimer and (B) aniline tetramer. The molecular weight for aniline trimer and tetramer is 291.0 and 366.3 Da, respectively.




Fig. S2 Optical microscopy images of NIH-3T3 cells incubated with $80 \mu\text{g mL}^{-1}$ of aniline oligomers for 24 h. (A) Control cells, (B) dimer, (C) trimer, (D) tetramer.




Fig. S3 Optical microscopy images of NIH-3T3 cells incubated with $80 \mu\text{g mL}^{-1}$ of aniline oligomers for 48 h. (A) Control cells, (B) dimer, (C) trimer, (D) tetramer.




Fig. S4 Optical microscopy images of A549 cells incubated with $80 \mu\text{g mL}^{-1}$ of aniline oligomers for 48 h. (A) Control cells, (B) dimer, (C) trimer, (D) tetramer.




Fig. S5 Optical microscopy images of A549 cells incubated with $80 \mu\text{g mL}^{-1}$ of aniline oligomers for 48 h. (A) Control cells, (B) dimer, (C) trimer, (D) tetramer.